

Appl. No. 09/601,004
Amdt. dated October 31, 2005
Reply to Office action of June 30, 2005

In the Claims:

Claims 1, 2, 4 are amended herein. Claims 3, 5, 7 and 8 are canceled. The remaining claims are not amended.

1. (currently amended) A two variable data interpolation system for processing image data, wherein a an image value between a plurality of discrete image data values is interpolated by performing convolution operation corresponding to the plurality of discrete data positioned at equal intervals in a two dimensional space using a sampling function that is differentiable finite times and has values of a local support,

wherein, with letting a third order B spline function be $F(t)$, the sampling function, $(H(t))$, is defined as follows:

$$H(t) = -F(t + 1/2)/4 + F(t) - F(t - 1/2)/4.$$

2. (currently amended) The two variable data interpolation system according to claim 1, wherein the sampling function is a function that ~~can be differentiated~~ is differentiable only once over a whole region.

3. (canceled)

4. (currently amended) The two variable data interpolation system according to claim 3 1, wherein the third order B spline function $F(t)$ is expressed as follows:

$$(4t^2 + 12t + 9)/4 \quad ; \quad -3/2 \leq t < -1/2$$

Appl. No. 09/601,004
 Amdt. dated October 31, 2005
 Reply to Office action of June 30, 2005

$$-2t^2 + 3/2 \quad ; \quad -1/2 \leq t < 1/2$$

$$(4t^2 - 12t + 9)/4 \quad ; \quad 1/2 \leq t < 3/2.$$

5. (canceled)

6. (previously presented) A two variable data interpolation system, wherein a value between a plurality of discrete data is interpolated by performing convolution operation corresponding to the plurality of discrete data positioned at equal intervals in a two dimensional space using a sampling function that is differentiable finite times and has values of a local support,

wherein, with letting a third order B spline function be $F(t)$, the sampling function, $H(t)$, is defined as follows:

$$H(t) = -F(t + 1/2)/4 + F(t) - F(t - 1/2)/4, \text{ comprising:}$$

discrete data extracting unit for extracting a plurality of discrete data that exist within a predetermined range around a data interpolating position that becomes an object of interpolation operation;

sampling function operating unit for calculating a value of the sampling function $H(t)$ for each of a plurality of discrete data extracted in this manner, with letting distance between the data interpolating position and discrete data be a t ; and

convolution operating unit for obtaining a value of the data interpolating position by performing convolution operation through adding values of the sampling function that are

Appl. No. 09/601,004
Amdt. dated October 31, 2005
Reply to Office action of June 30, 2005

calculated by the sampling function operating unit and correspond to the plurality of discrete data respectively.

7. (canceled)

8. (canceled)